



Published by DiscoverSys

Anti-inflammatory effect of red piper crocatum leaves extract decrease TNF- α and IL-6 levels in wistar rat with atherosclerosis



CrossMark

Sri Wahjuni,^{1*} Wita, I W,² Mantik Astawa, I N³

ABSTRACT

Background: This research aims to find a cure for anti-inflammation, based on the utilization of red piper *crocatum*. The research was started with descriptive study to explore active components of red piper *crocatum* leaf and followed by experimental study to investigate red piper *crocatum* activity of the leaf extract in anti-inflammation induced Wistar rat. In this research observed three dominant components: caryophyllene bicyclo [5.2.0] none, 2 methylene-4,8,8-trimethyl-4-vinyl; phytol; 5,9-propano-5H- benzocycloheptene, 6,7,8,9-tetrahydro-7,11-bis(methylene); 4,4-ethnedioxy-2-hexadecen-15-15 olide 1,4,9-trioxaspiro [4,15] eic os-6-en-8-one, 10 methyl; 1H-1,2,4-triazole-5(H)-thione,4-allyl-3-(3-furyl); Benzofuran,2,3-dihydro-2-methyl-7-phenyl which are possibly active to inhibit anti-inflammation to atherosclerosis. Bad eating habits also can cause various health problems, such as obesity, dyslipidemia, inflammation to atherosclerosis. This study was conducted to investigate of red piper *crocatum* leaves extract as an anti-inflammation through decrease of biochemistry markers TNF- α and IL-6 levels.

Method: This is a true experimental with randomized pre-test and post-test control group design, using 50 Wistar rats that are divided

into 5 groups: control group using 0 mg/kg BW red piper *crocatum* leaves extract, treatment group 1 using 50 mg/kg BW red piper *crocatum* leaves extract, treatment group 2 using 100 mg/kg BW red piper *crocatum* leaves extract, treatment group 3 using 150 mg/kg BW red piper *crocatum* leaves extract, and treatment group 4 200mg/kg BW red piper *crocatum* leaves extract.

Results: It was observed that intake of 150 mg/BW red piper *crocatum* leaves extract results in the highest significance decrease of 45.63% of TNF- α levels from $(28.62 \pm 1.25$ to 15.56 ± 7.20 pg/mL) and a significance decrease of 15.42% of IL-6 level from $(134.64 \pm 1.98$ to 113.87 ± 4.30 pg/mL).

Conclusion: It can be concluded that intake of red piper *crocatum* leaves extract acts as anti-inflammation for Wistar rats with atherosclerosis through decrease of TNF- α and IL-6 levels. Further research is required to determine whether the application of red piper *crocatum* leaves extract on human will result in similar effects of anti-inflammation.

Keywords: Consumption habit, instant food, dyslipidemia, atherosclerosis, red piper *crocatum* leaves extract.

Cite This Article: Wahjuni, S., Wita, I., Astawa, I. 2016. Anti-inflammatory effect of red piper crocatum leaves extract decrease TNF- α and IL-6 levels in wistar rat with atherosclerosis. *Bali Medical Journal* 5(2): 240-244. DOI: [10.15562/bmj.v5i2.210](https://doi.org/10.15562/bmj.v5i2.210)

¹Faculty of Mathematics and Basic Science, Udayana University, Bali, Indonesia

²Faculty of Medicine, Udayana University, Bali, Indonesia

³Faculty of Veterinary Medicine, Udayana University, Bali, Indonesia

INTRODUCTION

Inflammation is a complex biological response of body tissues to harmful stimuli, such as pathogens, damaged cells or irritants and a protective response involving immune cell, molecular mediator and blood vessels has been known that there is a significant correlation between high lipid serum levels and incidents of atherosclerosis, a trigger of coronary heart disease. Coronary heart disease presents as a results of blood circulation disturbance and abnormality of cardiac electricity or other forms of arrhythmia.

This leads to unorganized myocardial contraction, blood flow obstruction, and blood flow regurgitation. All of these conditions resulted in the return of the blood flow on each contraction back to the heart (shunts), blood flow abnormality and may end up with heart failure.¹ Atherosclerosis is a slowly progressive disease, present in large to medium arterial muscle and elastic artery. The main sites of atherosclerosis are abdominal aorta,

coronary artery, popliteal artery, thorax descending aorta, internal carotid artery, and the circle of Willis. Risk factors, such as hypertension, chronic hypercholesterolemia, immune system disturbance, toxin and virus are also involved in the arterial endothelial wall destruction. This damage induces permeability changes of endothelial cells and leads to the increase of plasma constituents, such as lipoprotein, that can easily enter to artery wall. Damaging of these endothelial cells could also change the trombosistein of lumen artery property that can leads to adhesion of thrombocyte to the blood and induce inflammation. If this damaging process exists continually for a long time, it will be followed by continuous atherosclerosis and leads to the thickening of tunica intima and results in disturbance of blood flow on that site.² Managing eating habits is one way to overcome this condition. Decreased plasma

*Correspondence to: Sri Wahjuni
Faculty of Mathematics and Basic Science, Udayana University, Denpasar
sriwahjunimanuaba@gmail.com

Table 1 Antioxidant activity test of red piper crocatum leaves crude extract

Sample	Time (minutes)	Test	Absorbance				% inhibition
			497 nm	517 nm	537 nm	A 517 nm	
Crude extract	5	DPPH sample	0.714	0.785	0.698	0.0790	71.22 %
			0.635	0.593	0.515	0.0180	
	60	DPPH sample	0.651	0.704	0.613	0.0720	85.42 %
			0.527	0.508	0.468	0.0105	

Table 2 Phytochemical Test of n-Butanol Extract

No.	Compounds	Reagent	Colour Changes	Results
1.	Alkaloid	Meyer	Yellow-orange (without white precipitate)	-
		Wagner	Yellow-brown (without brown precipitate)	-
2	Flavonoid	Willstatter	Yellow-crimson	+
		NaOH 10%	Yellow-brown	+
		H2SO4 concentrated	Yellow-crimson	+
		Bate Smith-Metcalf	Yellow-red	+
3	Triterpenoid	Lieberman-Burchard	Yellow-brown	+
		H2SO4 10%	Yellow-brown	+
4	Saponin	Hot water + HCl	No foam formation	-
5	Phenolate (Tannin)	Hot water + FeCl3	Yellow-greenish black	+
6	Steroid	Lieberman-Burchard	Yellow-brown	-
		H2SO2 10%	Yellow-brown	-

Remarks: (+) = containing tested compound
 (-) = not containing tested compound

Table 3 Compound Identified Based on GC-MS Chromatogram

Peaks	Retention time (tg)	% Area	Compound identified
1	14.06	1.39	Caryophyllene bicyclo[5.2.0] none,2 methylene-4,8,8-trimethyl-4-vinyl
2	21.55	4.67	Phytol
4	28.57	18.37	4,4-ethnedioxy-2-hexadecen-15-15 olide 1,4,9-trioxaspiro [4,15] eic os-6-en-8-one, 10 methyl
6	29.35	31.98	Benzofuran,2,3-dihydro-2-methyl-7-phenyl

Table 3 Mean of TNF- α serum levels data

Treatment	TNF- α (pg/mL)	
	Pre-test	Post-test
RPC 0 mg/kg BW (control)	28.98 \pm 6.00	28.11 \pm 5.94
RPC 50 mg/ kg BW	29.12 \pm 5.79	27.32 \pm 5.01
RPC 100 mg/kg BW	29.02 \pm 5.34	24.42 \pm 5.74
RPC 150 mg/kg BW	28.62 \pm 4.72	15.56 \pm 7.20
RPC 200 mg/ kg BW	29.02 \pm 5.06	26.02 \pm 8.34

RPC = Red piper crocatum

Table 4 Resume of Post Hoc Test of TNF- α Levels

Treatment	Mean Different of TNF- α (pg/mL)	p^*	
RPC 0 mg/ kg BW (control)	- RPC50 mg/ kg BW	0.79	0.770
	- RPC 100 mg/ kg BW	3.69	0.177
	- RPC 150 mg/ kg BW	12.55	0.001
	- RPC 200 mg/ kg BW	1.10	0.686
RPC 50 mg/ kg BW	- RPC 100 mg/ kg BW	2.90	0.287
	- RPC 150 mg/ kg BW	11.76	0.001
	- RPC 200 mg/ kg BW	0.30	0.911
RPC 100 mg/kg BW	-RPC 150mg/ kg BW	8.86	0.002
	- RPC 200mg/ kg BW	- 2.59	0.339
RPC 150 mg/ kg BW	- RPC 200 mg/ kg BW	- 1.45	0.001

RPC = Red piper crocatum
 * significant $p < 0.05$

performed based on post-test data and analyzed using one-way ANOVA. It was observed that there are differences between all treatments. The differences determined by using Post Hoc Test (LSD). The Post Hoc Test results are presented on [Table 4](#).

Data on [Table 3](#) were normally distributed with $p > 0.05$ and their variances were also homogenous with $p > 0.05$. The mean difference of various *Red piper crocatum* treatments can be obtained from the basis of post-test data, but only if all pre-test data are comparable. It was obtained that all pre-test data are comparable with $p < 0.05$, therefore, mean different of the treatment were obtained based on post-test data and analysed using one-way ANOVA.

There were significant differences of treatment obtained with $p < 0.0$. After that the data were analysed using Post Hoc Test to measure the difference. Post Hoc test results were summarized in [Table 6](#).

Decrease of IL-6 Levels

Mean of pre- and post-test data of IL-6 serum levels are presented on [Table 5](#).

Table 5 Mean of IL-6 Serum Levels Data Pre- and Post-Test

Treatment	IL-6 (pg/mL)	
	Pre-test	Post-test
RPC 0 mg/kg BW (control)	134.58 ± 2.21	133.15 ± 4.01
RPC 50 mg/kg BW	134.24 ± 2.64	130.28 ± 3.59
RPC 100 mg/kg BW	134.75 ± 2.51	127.20 ± 5.56
RPC 150 mg/kg BW	134.64 ± 1.98	113.87 ± 4.30
RPC 200 mg/kg BW	135.34 ± 4.57	120.87 ± 7.89

RPC = Red piper crocatum

Table 6 Resume of Post Hoc Test of IL-6 Levels

Treatment		Mean Different of IL-6 (rg/mL)	p*
RPC 0 mg/kg BW (control)	- RPC 50 mg/kg BW	2.87	0.232
	- RPC100 mg/kg BW	5.95	0.016
	- RPC 250 mg/kg BW	19.28	0.001
	- RPC 200 mg/kg BW	12.28	0.001
RPC 50 mg/kg BW	- RPC 100 mg/kg BW	- 3.09	0.201
	- RPC 150 mg/kg BW	16.41	0.001
	- RPC 200 mg/kg BW	9.41	0.001
RPC 100 mg/kg BW	- RPC 150 mg/kg BW	13.33	0.001
	- RPC 200 mg/kg BW	6.33	0.011
RPC 150 mg/kg BW	- RPC 200 mg/kg BW	- 7.00	0.001

RPC = Red piper crocatum
*Significant p < 0.05.

DISCUSSIONS

The research results indicate that the highest decrease of 12.55 pg/mL of TNF- α was obtained from intake of 150 mg/kg BW RPC. Increase of RPC to 200 mg/kg BW could not improve the decrease of TNF- α levels. This condition indicates that concentration of 200 mg/kg BW RPC has already active compound flavonoid. Therefore, it could not decrease of TNF- α levels any further. A research found that there is no transcription of NF-K β , so that no further production of TNF- α because of the active compound.⁷

Inflammation is a response to tissue damage during vascularization. This response is followed by an important process, such as endothelial process. Endothelium is an important part of blood vein that plays an important role in atherosclerosis. Endothelium is a main target of mechanical and chemical damage due to dyslipidemia risk factor. Chronic, continuous, and prolonged dyslipidemia resulted in pro-inflammation response and prothrombic which are initially acute becomes chronic. This will be followed by infiltration of leukocyte cells, mainly monocyte cells to lower subendothelial tissue to form macrophage cells. These cells will destroy all remains of LDL-C, oxidized to form foam cells and later will cause the formation of atheroma.⁸

The last two decades' research obtained that red piper crocatum leaves is effective as an anti-inflammation. This is because red piper crocatum is rich in flavonoids. These flavonoids are anti-inflammation flavonoids that can inhibit the pro-inflammatory cytokine.^{5,6,9} In this study, red piper crocatum leaves extract which is rich of flavanol was applied and proven to have an anti-inflammation effect. This anti-inflammation effect is due to activation of endothelial nuclear factor-kappa beta (ENF-K β) on peripheral vein. ENF-K β is a transcription factor distributed on all endothelial cells that has a role in controlling vascularization.

A research has found that the role of flavonoid as an anti-inflammation compound is due to their action as immunomodulator.⁹ In addition, their role as anti-inflammation compound is as a result of the flavanol effect. These acids are substrate for triggering the formation of pinocembrin and pinostrobin. These two flavanol are endothelium-dependent vasodilator, which can cause relaxation of ordinary coronary artery and paradoxical vasoconstriction on atherosclerosis artery.

Data in Table 4 indicate that there is a decrease of IL-6 levels since the intake of 50 mg/kg BW of RPC. Even though there is a decrease of IL-6 levels caused by treatment of 50 mg/kg BW RPC which is about 2.87 pg/mL, the decrease is not significant statistically with p > 0.05. Meanwhile an intake of 100 mg/kg BW RPC resulted in significant decrease of IL-6 levels by 5.95 pg/mL with p < 0.05. In addition, intake of 150 mg/BW RPC has also a similar trend to significantly decrease the IL-6 levels by 19.28 pg/mL with p < 0.05. However, increase of concentration to 200 mg/BW RPC intake did not significantly decrease the IL-6 levels, indicated by p > 0.05.

Red piper crocatum leaves are rich of pinocembrin and pinostrobin. These two flavanol are antioxidants that have anti-inflammation properties. In the endothelial cells experiencing activated inflammation, increased selectin and VCAM-1 expression were observed. VCAM-1 induces monocyte adhesion. This adhesion was also induced by pro-inflammation cytokines, such as IL-1 β and TNF- α . These cytokines were induced by CRP protein produced as a results of IL-6 response by signalling protease-activated receptor, uptake of oxLDL through oxLDL receptor-1 (LOX-1) and by interaction of CD40-CD40 ligand in artery intima.¹⁰ IL-6 has an important role in inflammation response and this cytokine is secreted by activated-macrophage, which leads to phebic and known as endogenous pyrogen. IL-6 was also initiated the acute phase response, marked by production of acute phase protein by the hepatocyte.¹¹

CONCLUSION

1. Intake of 150 mg/kg BW RPC decreases the TNF- α serum levels in Wistar rat with atherosclerosis by around 45.63% (from 28.62 ± 4.72 to 15.56 ± 7.20 pg/mL).
2. Intake of 0 mg/kg BW RPC decreases the IL-6 serum levels in Wistar rat with atherosclerosis by around 15.42% (from 134.64 ± 1.98 to 113.87 ± 4.30 pg/mL).

ACKNOWLEDGMENT

The author would like to thank Prof. Dr. dr. I Wayan Wita, SpJP. and Prof. Drh. N. Mantik Astawa, Ph.D. for their invaluable supervision on the Journal.

REFERENCES

1. Vinay, K., Abbas, A.K., and Nelson, F. 2004. Pathologic basic of disease 7th ed. Elsevier saunders. New York.
2. Szmítko, P.E., Wang, C.H., Weisel, R.D., De-Almmeida, J.R., Anderson, T.J., and Verma, S. 2003. New Markers of Inflammation and Endothelial Cell Activation. *Circulation* 108: 1917-1923.
3. Kanjwal, M. 2004. Peripheral Aterial Disease-The Silent Killer. *JK-Practitioner*, October-December:1 (no.4):235-232.
4. Stefan, J., Mikko, P.S.A., Bengt, K., and Jan-Nilsson. 1996. Human Monocytes/ Macrophages Release TNF- α in Response to Ox-LDL. *Arteriosclerosis, Thrombosis, and Vascular Biology*.16:1573-1579.
5. Ahmed, E. 2001. Immune Mechanism in Atherosclerosis. Dissertation. ISSN: 91-628-4612-4. Konferensrummet, Centrum för Molekular Medicin, Karolinska Sjukhuset.
6. Han, S.N., Leka, L.S., Lichtenstein, A.H., Ausman, L.M., Schaefer, E.J., and Meydani, S.N. 2002. Effect of Hydrogenated and Saturated, Relative to Polyunsaturated, Fat on Immune and Inflammatory Responses of Adults with Moderate Hypercholesterolemia. *Journal of Lipid Research*. 43(3): p. 445-52.
7. Chen, G., and Goeddel, D.V. 2002. TNFR-1 Signaling: A Beautiful Pathway. *Science*. 296: p. 1634 – 1635.
8. Baraas-Faisa. 2006. Kardiologi Molekuler Radikal Bebas, Disfungsi Endotel, Aterosklerosis. Antioksidan Latihan Fisik dan Rehabilitasi Jantung. Bagian Kardiologi FKUI/RS Jantung Harapan Kita. Jakarta.
9. Akhlaghi M., Brain B., 2009. Mechanism of flavonoid protection against myocardial ischemia reperfusion injury. *Journal of Molecular and Cellular Cardiology*, 46: 309-17.
10. Bonetti, P.O., Lerman, L.O., and Lerman, A. 2003. Endothelial Dysfunction: A Marker of Atherosclerosis Risk. *Atheroscler Thromb Vase Biol*. 23: p. 168-175.
11. Coico, R., Li, S.J., and Benyamin, E. 2003. Element of innate and acquired immunity *Immunology*. In: immunology. A Short Course, 5th ed. New Jersey, John Wiley & Sons p: 11 – 26.



This work is licensed under a Creative Commons Attribution